

Providence St. Joseph Health Providence St. Joseph Health Digital Commons

Journal Articles and Abstracts

7-24-2018

The Course of the V2 Segment of the Vertebral Arteries in Klippel-Feil Syndrome: A Case Report.

Asad Rizvi

Joe Iwanaga

Rod J Oskouian

Neurosurgery, Swedish Neuroscience Institute, Seattle, WA, USA.

Marios Loukas

R Shane Tubbs

Follow this and additional works at: <https://digitalcommons.psjhealth.org/publications>



Part of the [Neurology Commons](#)

Recommended Citation

Rizvi, Asad; Iwanaga, Joe; Oskouian, Rod J; Loukas, Marios; and Tubbs, R Shane, "The Course of the V2 Segment of the Vertebral Arteries in Klippel-Feil Syndrome: A Case Report." (2018). *Journal Articles and Abstracts*. 681.

<https://digitalcommons.psjhealth.org/publications/681>

This Article is brought to you for free and open access by Providence St. Joseph Health Digital Commons. It has been accepted for inclusion in Journal Articles and Abstracts by an authorized administrator of Providence St. Joseph Health Digital Commons. For more information, please contact digitalcommons@providence.org.

The Course of the V2 Segment of the Vertebral Arteries in Klippel-Feil Syndrome: A Case Report

Asad Rizvi ¹, Joe Iwanaga ², Rod J. Oskouian ³, Marios Loukas ⁴, R. Shane Tubbs ⁵

1. St. Georges University School of Medicine, St. Georges, GRD 2. Seattle Science Foundation, Seattle, USA 3. Neurosurgery, Swedish Neuroscience Institute, Seattle, USA 4. Department of Anatomical Sciences, St. George's University, School of Medicine, St. George's, GRD 5. Neurosurgery, Seattle Science Foundation, Seattle, USA

✉ **Corresponding author:** Joe Iwanaga, joei@seattlesciencefoundation.org
Disclosures can be found in Additional Information at the end of the article

Abstract

Klippel-Feil syndrome is a congenital disorder characterized by the fusion of one or more cervical vertebrae leading to limitations in the rotation, extension, and flexion of the neck and possible neurological symptoms. Other abnormalities can also be found in these patients. The anatomy of the vasculature can be abnormal in these patients including variations in the course and origin of the vertebral arteries potentially leading to intra-operative complications. Herein, we report a case of Klippel-Feil syndrome and detail the course of the vertebral arteries in an osteological specimen.

Categories: Pathology, Orthopedics

Keywords: klippel-feil syndrome, vertebral artery, ct angiography, imaging, cervical vertebrae, surgical management

Introduction

Klippel-Feil syndrome is characterized by the fusion of one or more cervical vertebrae [1]. Klippel and Feil [2] first identified this condition in a 46-year-old tailor having a short neck with a limited range of motion and a low posterior hairline. This condition can be inherited in autosomal dominant and recessive patterns, and many classification schemes have been proposed. Patients usually present with neurological symptoms, pain, and limited rotation, flexion, and extension of the neck. The age of presentation depends on the degree of severity of the fusion and other abnormalities, but these patients usually present at a later age [1,3]. Radiographs are essential in the evaluation of these patients to assess the degree of cervical fusion and other associated abnormalities [4]. Many management options exist depending on the symptoms and severity of the condition including modification of activities, bracing, arthrodesis, and placement of screws and plates [1].

The course of the vertebral arteries in Klippel-Feil syndrome is important, as an anomalous artery can be injured during surgical screw placement. The literature describes several vascular abnormalities associated with Klippel-Feil syndrome [5-8] including variations in the course and origin of the vertebral arteries [9-12]. Herein, we report a case of Klippel-Feil syndrome and detail an unusual course of the vertebral arteries in a bone specimen.

Case Presentation

How to cite this article

Rizvi A, Iwanaga J, Oskouian R J, et al. (July 24, 2018) The Course of the V2 Segment of the Vertebral Arteries in Klippel-Feil Syndrome: A Case Report. Cureus 10(7): e3038. DOI 10.7759/cureus.3038

Received 07/05/2018

Review began 07/10/2018

Review ended 07/13/2018

Published 07/24/2018

© Copyright 2018

Rizvi et al. This is an open access article distributed under the terms of the Creative Commons Attribution License CC-BY 3.0., which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Fusion of the C2 and C3 vertebrae was identified in a skeletal sample of an adult male (Figure 1). No other bony variations, such as cervical ribs, were noted in the specimen. In this specimen, irregularity of the right-sided superior articular facet surface and a short odontoid process and anomalous articular facet on the left side of the lateral odontoid process, characteristic of Klippel-Feil syndrome, were noted as well. The vertebral canal between C2 and C3 was stenotic, and the spinous processes of these vertebrae were bifid. There was a circumferential fusion between the two vertebrae. Upon passing red tubing through the adjacent transverse foramina of the left and right sides, the course of the vertebral arteries was determined to be very asymmetric (Figure 1). In the sagittal plane, the left vertebral artery traveled almost vertically but was angled posteriorly by about 10 degrees. However, the right vertebral artery at this vertebral level traveled more horizontally and was tilted posteriorly by about 45 degrees. Using the odontoid process as the midline, the left vertebral artery was found to be 2.8 cm from the midline and the right vertebral artery 2.5 cm from the midline.




FIGURE 1: First row: left, anterior view; middle, posterior view; right, right lateral view. Second row: left, left lateral view; middle, superior view; right, inferior view.

For all images, note the varied course of the V2 segment of the vertebral artery as demonstrated by the red tubing.

Discussion

Embryology of the cervical vertebrae and the vertebral artery

The vertebrae develop from the paraxial mesoderm formed by the process of gastrulation beginning on day 14 after conception. The paraxial mesoderm subdivides into smaller structures called somites through the process of segmentation between days 20 and 30. Somites further divide into sclerotomes (later form the vertebrae), myotomes, and dermatomes. A re-segmentation process occurs when the caudal part of the sclerotome of one somite fuses with the rostral part of the other, forming the vertebral body. Klippel-Feil syndrome is believed to result from mutations in the genes involved in regulating the segmentation and re-segmentation processes, and reports in the literature have identified several candidate genes [1].

The formation of the vascular system and the above-described development of the vertebrae occur simultaneously. During days 21 to 36 of gestation, six pairs of aortic arches are formed. The fourth right aortic arch forms the proximal part of the right subclavian artery between days 32 and 36, and the left fourth aortic arch persists to form a part of the adult aortic arch. During days 21 to 29, 30 pairs of dorsal intersegmental arteries arise from the dorsal aorta to supply the somites, and the neural tube and the first nine pairs anastomose with one another, forming the vertebral arteries. The proximal parts of the first six intersegmental arteries regress with the descent of the heart, leading to the vertebral arteries arising from the subclavian artery in adults. Finally, the seventh intersegmental artery forms the left and most of the right subclavian arteries. Bavinck et al. proposed that interruption in the blood supply through the vertebral arteries during embryonic development leads to Klippel-Feil syndrome and the associated vascular anomalies [13].

Klippel-Feil syndrome and the course of the vertebral artery

The usual course of the vertebral arteries can be divided into four parts: The first or the pre-foraminal part (V1) begins as the vertebral artery originates from the proximal subclavian artery and courses vertically upward to enter the transverse foramen of C6. In the second or the foraminal part (V2), it passes through the transverse foramen of C6-C1. The third or the extradural part (V3) courses on the upper surface of the posterior arch of the atlas (groove for vertebral arteries) and then enters the foramen magnum. The fourth or the intradural part (V4) is when the artery enters the dura mater [14].

Reports of variations in the course and origin of the vertebral arteries associated with Klippel-Feil syndrome include a case of a 15-year-old woman with the right vertebral artery originating from the carotid artery and then ascending vertically without entering the transverse foramina. Additionally, the left vertebral artery was hypoplastic in this patient [9]. Futane et al. [10] reported a small caliber right vertebral artery and a persistent left first intersegmental artery in a 26-year-old man with Klippel-Feil syndrome. Tian et al. [11] described a case of a 12-year-old female patient with Klippel-Feil syndrome with a high-riding right vertebral artery at C2. Another report [12] described a sinuous course of the V3 segment between C1 and C2 due to the ponticulus posticus of C2 in a 68-year-old woman with Klippel-Feil syndrome.

The management of patients with Klippel-Feil syndrome depends on the severity of the condition and can involve surgical screw placement. The vertebral artery is at risk of injury during such procedures [15-16]. Therefore, pre-operative imaging of the craniocervical junction with, for example, computed tomography (CT) angiography is essential for identifying the course of the vertebral arteries to avoid intra-operative complications [10,17].

Conclusions

The literature describes some variations in the course and origin of the vertebral arteries in Klippel-Feil syndrome. These reports underscore the importance of pre-surgical or intraoperative imaging to identify any anomalies in the vertebral arteries and the need to exercise prudence during surgery to avoid potential complications. The case presented here offers a unique window into the anomalous course of the vertebral arteries in a skeletal specimen of an adult male with Klippel-Feil syndrome.

Additional Information

Disclosures

Human subjects: All authors have confirmed that this study did not involve human participants or tissue. **Conflicts of interest:** In compliance with the ICMJE uniform disclosure form, all authors declare the following: **Payment/services info:** All authors have declared that

no financial support was received from any organization for the submitted work. **Financial relationships:** All authors have declared that they have no financial relationships at present or within the previous three years with any organizations that might have an interest in the submitted work. **Other relationships:** All authors have declared that there are no other relationships or activities that could appear to have influenced the submitted work.

References

1. Tracy MR, Dormans JP, Kusumi K: Klippel-Feil syndrome: clinical features and current understanding of etiology. *Clin Orthop Relat Res*. 2004, 424:183-90. [10.1097/01.blo.0000130267.49895.20](https://doi.org/10.1097/01.blo.0000130267.49895.20)
2. Klippel M, Feil A: The classic: A case of absence of cervical vertebrae with the thoracic cage rising to the base of the cranium (cervical thoracic cage). *Clin Orthop Relat Res*. 1975, 109:3-8. [10.1097/00003086-197506000-00002](https://doi.org/10.1097/00003086-197506000-00002)
3. Dietz F: Congenital Abnormalities of the Cervical Spine . *The Pediatric Spine—Principles and Practice*. Weinstein SL (ed): Lippincott Williams & Wilkins, Philadelphia; 2001. 2:239-251.
4. Manaligod JM, Bauman NM, Menezes AH, Smith RJH: Cervical vertebral anomalies in patients with anomalies of the head and neck. *Ann Otol Rhinol Laryngol*. 1999, 108:925-933. [10.1177/000348949910801002](https://doi.org/10.1177/000348949910801002)
5. Paksoy Y, Seker M, Kalkan E: Klippel-Feil syndrome associated with persistent trigeminal artery. *Spine*. 2004, 29:193-6. [10.1097/00007632-200405010-00024](https://doi.org/10.1097/00007632-200405010-00024)
6. Franzen D, Schulte B, Beyer D, Nidel J, Koebke J, de Vivie R: Klippel-Feil syndrome associated with aortic coarctation. *Cardiovasc Pathol*. 2003, 12:115-7. [10.1016/s1054-8807\(02\)00187-4](https://doi.org/10.1016/s1054-8807(02)00187-4)
7. Rielly JP: Intrathoracic carotid bifurcation in Klippel-Feil syndrome. *J Vasc Surg*. 2008, 47:1071-3. [10.1016/j.jvs.2007.11.022](https://doi.org/10.1016/j.jvs.2007.11.022)
8. Brill CB, Peyster RG, Keller MS, Galtman L, Opitz JM, Reynolds JF: Isolation of the right subclavian artery with subclavian steal in a child with Klippel-Feil anomaly: an example of the subclavian artery supply disruption sequence. *Am J Med Genet*. 1987, 26:933-40. [10.1002/ajmg.1320260424](https://doi.org/10.1002/ajmg.1320260424)
9. Ulusoy OL, Sasani H, Barlas SB, Mutlu A, Sasani M: A case of anomalous origin and course of vertebral artery in a patient with Klippel Feil syndrome. *Korean J Radiol*. 2016, 17:554-557. [10.3348/kjr.2016.17.4.55](https://doi.org/10.3348/kjr.2016.17.4.55)
10. Futane S, Salunke P: Klippel-Feil syndrome with atlanto-axial dislocation, anomalous vertebral artery, dextrocardia and situs inversus. *Clin Neurol Neurosurg*. 2013, 115:2304-2306. [10.1016/j.clineuro.2013.08.011](https://doi.org/10.1016/j.clineuro.2013.08.011)
11. Tian Y, Fan D, Xu N, Wang S: “Sandwich Deformity” in Klippel-Feil syndrome: a “Full-Spectrum” presentation of associated craniovertebral junction abnormalities. *J Clin Neurosci*. 2018, 53:247-249. [10.1016/j.jocn.2018.04.047](https://doi.org/10.1016/j.jocn.2018.04.047)
12. Lampropoulou-Adamidou K, Athanassacopoulos M, Karampinas PK, Vlamis J, Korres DS, Pneumáticos SG: Congenital variations of the upper cervical spine and their importance in preoperative diagnosis. A case report and a review of the literature. *Eur J Orthop Surg Traumatol*. 2013, 23:101-105. [10.1007/s00590-013-1216-z](https://doi.org/10.1007/s00590-013-1216-z)
13. Bavinck JNB, Weaver DD: Subclavian artery supply disruption sequence: hypothesis of a vascular etiology for Poland, Klippel-Feil, and mobius anomalies. *Am J of Med Genet*. 1986, 23:903-918. [10.1002/ajmg.1320230405](https://doi.org/10.1002/ajmg.1320230405)
14. Standring S: *Gray's Anatomy: The Anatomical Basis of Clinical Practice*. Churchill Livingstone, London; 2008.
15. Young JP, Young PH, Ackermann MJ, Anderson PA, Riew KD: The ponticulus posticus: implications for screw insertion into the first cervical lateral mass. *J Bone Joint Surg Am*. 2005, 87:2495-2498. [10.2106/00004623-200511000-00017](https://doi.org/10.2106/00004623-200511000-00017)
16. Tardieu GG, Edwards B, Alonso F, et al.: Aortic arch origin of the left vertebral artery: an anatomical and radiological study with significance for avoiding complications with anterior approaches to the cervical spine. *Clin Anat*. 2017, 30:811-816. [10.1002/ca.22923](https://doi.org/10.1002/ca.22923)
17. Yamazaki M, Okawa A, Furuya T, et al.: Anomalous vertebral arteries in the extra- and intraosseous regions of the craniovertebral junction visualized by 3-dimensional computed tomographic angiography: analysis of 100 consecutive surgical cases and review of the

literature. Spine. 2012, 37:1389-97. [10.1097/brs.0b013e31826a0c9f](https://doi.org/10.1097/brs.0b013e31826a0c9f)